
Astragalus diaphanus var. diurnus

Fabaceae South John Day milkvetch

Gerald D. Carr






Aaron Liston



Gerald D. Carr








Prostrate annual or biennial arising from a slender taproot. Stems 3-10, rarely numerous, 15-40 cm high, red tinged, strigulose. Leaves 2-5 (6) cm long, often red-tinged; short, slender petioles; 7-15 leaflets, oblong or obovate, 3-12 mm long, glabrous above, strigulose beneath. Inflorescence a raceme; loosely (3) 7-15 flowered; peduncles 1-5.5 cm long, mostly shorter than the leaves. Flowers 4-9 mm long; petals whitish; banner lavender-veined, 7-9 mm long; wings 6-8 mm long; keel purple-tipped, 4-6 mm long; calyx stigulose with black or black and white hairs, 3.5-4.5 mm long. Pods ovoid, distinctly inflated, 1-3 cm long, 3-9 mm in diameter, sulcate dorsally, purple-green, becoming translucent; intruded sutures forming a scarcely developed partition.

Lookalikes Astragalus diaphanus differs from featured plant by its fruits that are strongly curved and not inflated

Astragalus diaphanus Douglas ex Hook. var. diurnus (S. Watson) Barneby ex M. Peck

South John Day milkvetch PLANTS symbol: ASDI2

August 2019 status Oregon:LT; ORBIC: List 1



Distribution: Valley of the South John Day River, Grant Co.

Habitat: Shallow gravely soil over basalt, sandbars or sandy banks of intermittent rivers.

010.

Elevation: 700-1400 m

Best survey time (in flower): mid-April - June

Associated species:

Artemisia tridentata (Big sagebrush)
Artemisia rigida (Scabland sagebrush)
Eriogonum compositum (Arrowleaf buckwheat)
Eriogonum sphaerocephalum (Rock buckwheat)
Eriophyllum lanatum (Common woolly sunflower)
Juniperus occidentalis (Western juniper)
Ericameria nauseosa (Rubber rabbitbrush)
Lewisia rediviva (Bitterroot)
Salvia dorrii (Purple sage)





