

ten scapelike, 15-25 cm long. Bulbs whitish, coated with a dull brown membranous coat. Leaf basal, solitary, erect, to 30 cm long, 3-7 mm wide, inner surface densely hairy, outer surface glabrous. Inflorescence 1-7 flowered on erect pedicel; bracts 1 to several, 2.5-3 cm long. Flowers white with reddish striations from base to gland, a broad lavender chevron extending to petal edges just above the gland; gland deeply depressed, base green; yellow hairs surround the gland that grade to white at petal apex; sepals ovate-acuminate, about 20 mm long, 8 mm wide; inner surface of petals densely hairy, outer surface glabrous, edges ciliate; filaments about 7 mm long; anthers reddish-brown, 3-7 mm long. Fruit a capsule, straw colored; nodding, elliptic-elongate, 3 winged, 3-4 cm long. Seeds light straw yellow with a rough surface.

Lookalikes Calochortus tolmiei Calochortus greenei

differs from featured plant by


its densely hairy petals with pink to purple striations. its lilac petals and glabrous inner leaf surface and earlier bloom time.



Gerald D. Carr










Calochortus coxii M.R. Godfrey & Callahan

Cox's mariposa lily PLANTS symbol: CACO41 *August 2019 status* Federal:SOC; Oregon:LE; ORBIC: List 1



Distribution: From the Umpqua River to Myrtle Creek Mountain, Douglas Co.

Habitat: Serpentine soils on north facing, open, grassy or wooded slopes.

Elevation: 200-800 m

Best survey time (in flower): late June - early July

Associated species:

Pinus jeffreyi (Jeffrey pine) Calocedrus decurrens (Incense cedar) Danthonia californica (California danthonia) Aspidotis densa (Podfern)

