

Hackelia cronquistii

Boraginaceae

Cronquist's stickseed

Robert L. Carr




Robert L. Carr



Robert L. Carr



Robert L. Carr



Illustrated by Frank A. Lang
From: Threatened and Endangered
Vascular Plants of Oregon (1982)
R. J. Meinke

Erect perennial herb from a taproot and compactly branched caudex. Stems glabrous below, sparsely strigose above, 20-65 cm tall. Leaves hispid to strigose; basal leaves long petiolate, narrowly elliptic to lance-elliptic, acute, 6-21 cm long, 5-35 mm wide; cauline leaves usually sessile, narrowly elliptic to narrowly-oblong, 2.5-14.5 cm long, 2-13 mm wide, narrowly acute, progressively smaller upward. Inflorescence panicate; pedicels 8-14 mm long; bracts small and inconspicuous. Calyx lobes 1.7-2.8 mm long oblong-lanceolate to lanceolate. Corolla limb white, tinged with blue, 8-15 mm wide, tube 2-2.2 mm long; fornications papillate or papillate puberulent. Fruit a nutlet, 2.5-3.5 mm long, lanceolate to lance-ovate.

Lookalikes
Hackelia spp.

differs from featured plant by
having hairy stems throughout and blue or white corollas with yellow eyes.

best survey times
J | F | M | A | M | J | J | A | S | O | N | D




Hackelia cronquistii J.L. Gentry

Cronquist's stickseed

PLANTS symbol: HACR4

August 2019 status Oregon:LT; ORBIC: List 1



Distribution: Northeastern Malheur Co. and extreme southeastern Baker Co.

Habitat: Sand and sandy loam.

Elevation: 600–1100 m


Best survey time (in flower): Late April – May

Associated species:

Artemisia tridentata (Big sagebrush)

Eriophyllum lanatum var. *integrifolium* (Woolly sunflower)

Balsamorhiza sagittata (Arrowleaf balsamroot)



Oregon Flora

<https://oregonflora.org>

Published August 2019